

# Generalized Quadrangles of Order $(s, s^2)$ , I

J. A. THAS

*Department of Pure Mathematics and Computer Algebra, University of Gent,  
Krijgslaan 281, B-9000 Gent, Belgium*

*Communicated by Francis Buekenhout*

Received January 5, 1992

In this paper generalized quadrangles of order  $(s, s^2)$ ,  $s > 1$ , satisfying property (G) at a line, at a pair of points, or at a flag, are studied. Property (G) was introduced by S. E. Payne (*Geom. Dedicata* 32 (1989), 93–118) and is weaker than 3-regularity (see S. E. Payne and J. A. Thas, “Finite Generalized Quadrangles,” Pitman, London, 1984). It was shown by Payne that each generalized quadrangle of order  $(s^2, s)$ ,  $s > 1$ , arising from a flock of a quadratic cone, has property (G) at its point  $(\infty)$ . In particular translation generalized quadrangles satisfying property (G) are considered here. As an application it is proved that the Roman generalized quadrangles of Payne contain at least  $s^3 + s^2$  classical subquadrangles  $Q(4, s)$ . Also, as a by-product, several classes of ovoids of  $Q(4, s)$ ,  $s$  odd, are obtained; one of these classes is new. The goal of Part II is the classification of all translation generalized quadrangles satisfying property (G) at some flag  $((\infty), L)$ . © 1994 Academic Press, Inc.

## 1. GENERALIZED QUADRANGLES

### 1.1. Introduction

A (finite) generalized quadrangle (GQ) is an incidence structure  $\mathcal{S} = (P, B, I)$  in which  $P$  and  $B$  are disjoint (nonempty) sets of objects called points and lines (respectively), and for which  $I$  is a symmetric point-line incidence relation satisfying the following axioms:

- (i) Each point is incident with  $1 + t$  lines ( $t \geq 1$ ) and two distinct points are incident with at most one line.
- (ii) Each line is incident with  $1 + s$  points ( $s \geq 1$ ) and two distinct lines are incident with at most one point.
- (iii) If  $x$  is a point and  $L$  is a line not incident with  $x$ , then there is a unique pair  $(y, M) \in P \times B$  for which  $x I M I y I L$ .

The integers  $s$  and  $t$  are the parameters of the GQ and  $\mathcal{S}$  is said to have order  $(s, t)$ ; if  $s = t$ ,  $\mathcal{S}$  is said to have order  $s$ . There is a point-line duality for GQ for which in any definition or theorem the words "point" and "line" are interchanged and the parameters  $s$  and  $t$  are interchanged. Normally, we assume without further notice that the dual of a given theorem or definition has also been given.

Given two (not necessarily distinct) points  $x, y$  of the GQ  $\mathcal{S}$ , we write  $x \sim y$  and say that  $x$  and  $y$  are collinear, provided that there is some line  $L$  for which  $x I L I y$ . And  $x \not\sim y$  means that  $x$  and  $y$  are not collinear. Dually, for  $L, M \in B$ , we write  $L \sim M$  or  $L \not\sim M$  according as  $L$  and  $M$  are concurrent or nonconcurrent, respectively. If  $x \sim y$  (resp.,  $L \sim M$ ) we may also say that  $x$  (resp.,  $L$ ) is orthogonal or perpendicular to  $y$  (resp.,  $M$ ). The line (resp., point) which is incident with distinct collinear points  $x, y$  (resp. distinct concurrent lines  $L, M$ ) is denoted by  $xy$  (resp.,  $LM$  or  $L \cap M$ ).

For  $x \in P$  put  $x^\perp = \{y \in P \mid y \sim x\}$ , and note that  $x \in x^\perp$ . The trace of a pair  $\{x, y\}$  of distinct points is defined to be the set  $x^\perp \cap y^\perp$  and is denoted  $\text{tr}\{x, y\}$  or  $\{x, y\}^\perp$ . We have  $|\{x, y\}^\perp| = s + 1$  or  $t + 1$  according as  $x \sim y$  or  $x \not\sim y$ . More generally, if  $A \subset P$ ,  $A$  "perp" is defined by  $A^\perp = \bigcap \{x^\perp \mid x \in A\}$ . For  $x \neq y$ , the span of the pair  $\{x, y\}$  is  $\text{sp}\{x, y\} = \{x, y\}^{\perp\perp} = \{u \in P \mid u \in z^\perp \forall z \in x^\perp \cap y^\perp\}$ . We have  $|\{x, y\}^{\perp\perp}| = s + 1$  or  $|\{x, y\}^{\perp\perp}| \leq t + 1$  according as  $x \sim y$  or  $x \not\sim y$ .

A triad (of points) is a triple of pairwise noncollinear points. Given a triad  $T = \{x, y, z\}$ , a center of  $T$  is just a point of  $T^\perp$ .

For definitions, notations, and results neither given herein nor cited explicitly, we refer to the monograph by Payne and Thas [11].

### 1.2. Restrictions on the Parameters

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(s, t)$ . If  $|P| = v$  and  $|B| = b$ , then  $v = (s + 1)(st + 1)$  and  $b = (t + 1)(st + 1)$ ; see [11, 1.2.1]. Also,  $s + t$  divides  $st(s + 1)(t + 1)$ ; see [11, 1.2.2].

If  $s > 1$  and  $t > 1$ , then  $t \leq s^2$ , and dually  $s \leq t^2$ ; see [11, 1.2.3]. Also, for  $s > 1$  and  $t > 1$ ,  $s^2 = t$  iff each triad (of points) has a constant number of centers, in which case this constant number of centers is  $s + 1$ ; see [11, 1.2.4].

## 2. REGULARITY, ANTIREGULARITY, 3-REGULARITY, AND PROPERTY (G)

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(s, t)$ .

### 2.1. Regularity

If  $x \sim y$ ,  $x \neq y$ , or if  $x \not\sim y$  and  $|\{x, y\}^{\perp\perp}| = t + 1$ , we say that the pair  $\{x, y\}$  is regular. The point  $x$  is regular provided  $\{x, y\}$  is regular for all

$y \in P$ ,  $y \neq x$ . A point  $x$  is coregular provided each line incident with  $x$  is regular.

If  $\mathcal{S}$  contains a regular pair  $\{x, y\}$ , then either  $s=1$  or  $s \geq t$ ; see [11, 1.3.6]. A coregular point in a GQ of order  $s$ ,  $s > 1$ , is regular iff  $s$  is even; see [11, 1.5.2].

Let  $x$  be a regular point of  $\mathcal{S}$ . Then the incidence structure with pointset  $x^\perp - \{x\}$ , with lineset the set of spans  $\{y, z\}^{\perp\perp}$ , where  $y, z \in x^\perp - \{x\}$ ,  $y \not\sim z$ , and with the natural incidence, is the dual of a net (cf. [2]) of order  $s$  and degree  $t+1$ ; see [11, 1.3.1]. If in particular  $s=t>1$ , there arises a dual affine plane of order  $s$ . Also, in this case the incidence structure  $\pi_x$  with pointset  $x^\perp$ , with lineset the set of spans  $\{y, z\}^{\perp\perp}$ , where  $y, z \in x^\perp$ ,  $y \neq z$ , and with the natural incidence, is a projective plane of order  $s$ .

EXAMPLES. (a) Let  $Q(n, q)$  be the GQ arising from the nonsingular quadric  $Q$  of Witt index 2 in  $PG(n, q)$ ,  $n \in \{3, 4, 5\}$ . All lines of  $Q(n, q)$  are regular [11, 3.3.1]. All points of  $Q(3, q)$  are regular; all points of  $Q(4, q)$  are regular iff  $q$  is even; and no point of  $Q(5, q)$  is regular [11, 3.3.1].

(b) Let  $H(n, q^2)$  be the GQ arising from the nonsingular Hermitian variety  $H$  of  $PG(n, q^2)$ ,  $n \in \{3, 4\}$ . All points of  $H(3, q^2)$  are regular and no line of  $H(3, q^2)$  is regular [11, 3.3.1]. No point and no line of  $H(4, q^2)$  is regular [11, 3.3.1].

(c) Since the GQ  $W(q)$  arising from a symplectic polarity  $\zeta$  of  $PG(3, q)$  is isomorphic to the dual of  $Q(4, q)$  (cf. [11, 3.2.1]), all points of  $W(q)$  are regular for all  $q$  and all lines of  $W(q)$  are regular iff  $q$  is even.

(d) Let  $T_2(O)$  be the GQ arising from an oval  $O$  of  $PG(2, q)$ . Then the point  $(\infty)$  is coregular for  $T_2(O)$  [11, 3.3.2].

Let  $T_3(O)$  be the GQ arising from an ovoid  $O$  of  $P(3, q)$ . Then the point  $(\infty)$  is coregular for  $T_3(O)$  [11, 3.3.2].

Other results on regularity in  $T_n(O)$ ,  $n \in \{2, 3\}$ , can be found in [11, 3.3.3].

(e) For any translation generalized quadrangle  $\mathcal{S}^{(p)}$  the base point  $p$  is coregular [11, 8.1 and 8.2].

(f) Let  $\mathcal{S}(F)$  be the GQ of order  $(q^2, q)$  arising from the flock  $F$  of the quadratic cone  $K$  of  $PG(3, q)$ ; see [16]. Then it is easy to check that the point  $(\infty)$  is regular.

**THEOREM 2.1.1.** *A GQ  $\mathcal{S}$  of order  $s$  ( $s > 1$ ) is isomorphic to  $W(s)$  iff all its points are regular.*

*Proof.* See [11, 5.2.1]. ■

## 2.2. Antiregularity

The pair  $\{x, y\}$ ,  $x \not\sim y$ , is antiregular provided  $|z^\perp \cap \{x, y\}^\perp| \leq 2$  for all  $z \in P - \{x, y\}$ . A point  $x$  is antiregular provided  $\{x, y\}$  is antiregular for all  $y \in P - x^\perp$ .

If  $\{x, y\}$  is antiregular with  $s = t$ , then  $s$  is odd; see [11, 1.5.1]. If  $1 < s < t$ , then  $\mathcal{S}$  has no antiregular pair of points; see [11, 1.3.6]. A coregular point in a GQ of order  $s$ ,  $s > 1$ , is antiregular iff  $s$  is odd; see [11, 1.5.2].

Let  $x$  be an antiregular point of the GQ  $\mathcal{S} = (P, B, I)$  of order  $s$ ,  $s \neq 1$ . Then the incidence structure with pointset  $x^\perp - \{x\}$ , with lineset the set of spans  $\{x, y\}^{\perp\perp}$ ,  $y \sim x$ , and  $y \neq x$ , with circleset the set of traces  $\{x, z\}^\perp$ ,  $z \not\sim x$ , and with the natural incidence, is a Laguerre plane  $\mathcal{L}$  of order  $s$ ; see [10]. For each point  $u$  of  $\mathcal{L}$  the derived or internal affine plane of  $\mathcal{L}$  at  $u$  will be denoted by  $\mathcal{L}_u$  or  $\pi(x, u)$ ; points of  $\pi(x, u)$  are the points of  $x^\perp$  not collinear with  $u$ , and the lines of  $\pi(x, u)$  are the spans  $\{x, y\}^{\perp\perp}$ , with  $y \sim x$  and  $y \not\sim u$ , and the traces  $\{x, z\}^\perp$ , with  $z \not\sim x$  and  $z \sim u$ .

EXAMPLES [11, 3.3.1]. All points of  $Q(4, q)$  are antiregular iff  $q$  is odd; no line of  $Q(4, q)$  is antiregular. Since  $W(q)$  is isomorphic to the dual of  $Q(4, q)$ , all lines of  $W(q)$  are antiregular iff  $q$  is odd, and no point of  $W(q)$  is antiregular.

THEOREM 2.2.1. *Let  $\mathcal{S}$  be a GQ of order  $s$ ,  $s \neq 1$ , having an antiregular point  $x$ . Then  $\mathcal{S}$  is isomorphic to  $Q(4, s)$  iff there is a point  $y$ ,  $y \in x^\perp - \{x\}$ , for which the associated affine plane  $\pi(x, y)$  is Desarguesian.*

*Proof.* See [11, 5.2.7]. ■

## 2.3. 3-Regularity

Let  $s^2 = t > 1$ , so that by 1.2 we have  $|\{x, y, z\}^\perp| = s + 1$  for any triad  $\{x, y, z\}$ . Evidently  $|\{x, y, z\}^{\perp\perp}| \leq s + 1$ . We say  $\{x, y, z\}$  is 3-regular provided  $|\{x, y, z\}^{\perp\perp}| = s + 1$ . The point  $x$  is called 3-regular iff each triad containing  $x$  is 3-regular.

Let  $\mathcal{S}$  be a GQ of order  $(s, s^2)$ ,  $s \neq 1$ , and suppose that any triad contained in  $\{x, y\}^\perp$ ,  $x \not\sim y$ , is 3-regular. Then the incidence structure with pointset  $\{x, y\}^\perp$ , with circleset the set of elements  $\{z, z', z''\}^{\perp\perp}$ , where  $z, z', z''$  are distinct points in  $\{x, y\}^\perp$  and, with the natural incidence, is an inversive plane of order  $s$ ; see [11, 1.3.3].

If  $\{x, y, z\} = T$  is a 3-regular triad of the GQ  $\mathcal{S} = (P, B, I)$  of order  $(s, s^2)$ ,  $s \neq 1$ , then each point of  $P - (T^\perp \cup T^{\perp\perp})$  is collinear with precisely two points of  $T^\perp \cup T^{\perp\perp}$ ; see [11, 1.4.1].

Let  $\{x, y, z\} = T$  be a 3-regular triad of the GQ  $\mathcal{S} = (P, B, I)$  of order  $(s, s^2)$ ,  $s > 1$ , and let  $P'$  be the set of all points incident with lines of the

form  $uv$ ,  $u \in T^\perp$ , and  $v \in T^{\perp\perp}$ . If  $L$  is a line which is incident with no point of  $T^\perp \cup T^{\perp\perp}$  and if  $k$  is the number of points in  $P'$  which are incident with  $L$ , then  $k \in \{0, 2\}$  if  $s$  is odd and  $k \in \{1, s+1\}$  if  $s$  is even; see [11, 2.6.1]. If  $s$  is even, if  $B'$  is the set of all lines in  $B$  which are incident with at least two points of  $P'$ , and if  $I'$  is the restriction of  $I$  to  $(P' \times B') \cup (B' \times P')$ , then  $\mathcal{S}' = (P', B', I')$  is a subquadrangle of  $\mathcal{S}$  of order  $s$ ; moreover  $\{x, y\}$  is a regular pair of  $\mathcal{S}'$ , with  $\{x, y\}^{\perp'} = \{x, y, z\}^\perp$  and  $\{x, y\}^{\perp'\perp'} = \{x, y, z\}^{\perp\perp}$  (see [11, 2.6.2]).

EXAMPLES. (a) All points of  $Q(5, q)$  are 3-regular [11, 3.3.1]. Since  $H(3, q^2)$  is isomorphic to the dual of  $Q(5, q)$ , all lines of  $H(3, q^2)$  are 3-regular.

(b) The point  $(\infty)$  of the GQ  $T_3(O)$  is 3-regular [11, 3.3.2]. If  $T_3(O)$  has a 3-regular point other than  $(\infty)$ , then  $O$  is an elliptic quadric and  $T_3(O)$  is isomorphic to  $Q(5, q)$  [11, 3.3.3].

THEOREM 2.3.1. (i) A GQ of order  $(s, s^2)$ ,  $s > 1$ , is isomorphic to  $T_3(O)$  iff it has a 3-regular point.

(ii) Let  $\mathcal{S}$  be a GQ of order  $(s, s^2)$ ,  $s > 1$ , with  $s$  odd. Then  $\mathcal{S} \cong Q(5, s)$  iff  $\mathcal{S}$  has a 3-regular point.

(iii) Let  $\mathcal{S}$  be a GQ of order  $(s, s^2)$  with  $s$  even. Then  $\mathcal{S} \cong Q(5, s)$  iff one of the following holds:

(a) All points of  $\mathcal{S}$  are 3-regular.

(b)  $\mathcal{S}$  has at least one 3-regular point not incident with some regular line.

*Proof.* See [11, 5.3.1 and 5.3.3]. ■

#### 2.4. Property (G) for GQ of Order $(s, s^2)$

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(s, s^2)$ ,  $s \neq 1$ . Let  $x_1, y_1$  be distinct collinear points. We say that the pair  $\{x_1, y_1\}$  has property (G) or that  $\mathcal{S}$  has property (G) at  $\{x_1, y_1\}$ , if every triad  $\{x_1, x_2, x_3\}$ , with  $y_1 \in \{x_1, x_2, x_3\}^\perp$ , is 3-regular. Then also every triad  $\{y_1, y_2, y_3\}$ , with  $x_1 \in \{y_1, y_2, y_3\}^\perp$ , is 3-regular. The GQ  $\mathcal{S}$  has property (G) at the line  $L$ , or the line  $L$  has property (G), if each pair of points  $\{x, y\}$ ,  $x \neq y$ , and  $x I L I y$ , has property (G). If  $(x, L)$  is a flag, that is, if  $x I L$ , then we say that  $\mathcal{S}$  has property (G) at  $(x, L)$ , or that  $(x, L)$  has property (G), if every pair  $\{x, y\}$ ,  $x \neq y$ , and  $y I L$ , has property (G).

The point  $x$  of the GQ  $\mathcal{S}$  is 3-regular iff each flag  $(x, L)$ ,  $x I L$ , has property (G).

If  $s$  is even and if  $\mathcal{S}$  contains a pair of points having property (G), then by 2.3  $\mathcal{S}$  contains subquadrangles of order  $s$ .

In [9] property (G) is defined for any GQ of order  $(s, t)$ ,  $t > s > 1$ .

EXAMPLE. If  $\mathcal{S}(F)$  is the GQ of order  $(q^2, q)$  arising from the flock  $F$  of the quadratic cone  $K$  of  $PG(3, q)$ , then by [9]  $\mathcal{S}$  has property (G) at the point  $(\infty)$ .

*Goal of the paper.* In the subsequent sections GQ of order  $(s, s^2)$ ,  $s \neq 1$ , having property (G) at some pair of points, or at some line, or at some flag, will be considered. In particular translation generalized quadrangles of order  $(s, s^2)$ ,  $s \neq 1$ , with base point  $(\infty)$  and having property (G) either at some flag  $((\infty), L)$  or at some line  $L$  incident with the point  $(\infty)$ , are studied in detail.

### 3. GENERALIZED QUADRANGLES OF ORDER $(q, q^2)$ , $q \neq 1$ , SATISFYING PROPERTY (G)

#### 3.1. The Affine Space $\mathcal{S}_{xy}$ (S. E. Payne and J. A. Thas)

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(q, q^2)$ ,  $q \neq 1$ , for which the pair of points  $\{x, y\}$ ,  $x \sim y$ , has property (G). Now we introduce the following incidence structure  $\mathcal{S}_{xy} = (P_{xy}, B_{xy}, I_{xy})$ :

- (i)  $P_{xy} = x^\perp - \{x, y\}^{\perp\perp}$ .
- (ii) Elements of  $B_{xy}$  are of two types: (a) the sets  $\{y, z, u\}^{\perp\perp} - \{y\}$ , with  $\{y, z, u\}$  a triad having  $x$  as center, and (b) the sets  $\{x, w\}^\perp - \{x\}$ , with  $x \sim w \not\sim y$ .
- (iii)  $I_{xy}$  is containment.

Then  $|P_{xy}| = q^3$ ,  $B_{xy}$  contains  $q^3(q^3 - q)/q(q - 1) = q^4 + q^3$  elements of type (a) and  $q^2$  elements of type (b).

LEMMA 3.1.1. Let  $M_1, M_2$  be different elements of type (a) of  $B_{xy}$ , let  $L_1, L_2$  be different elements of type (b) of  $B_{xy}$ , and assume that  $L_i \cap M_j \neq \emptyset$  for all  $i, j \in \{1, 2\}$ . Then each element of type (b) of  $B_{xy}$  which contains a point of  $M_1$  also contains a point of  $M_2$ .

*Proof.* Let  $L_i \cap M_j = \{x_{ij}\}$ . Since  $M_1 \neq M_2$ , we have  $\{x_{11}, x_{21}\} \neq \{x_{12}, x_{22}\}$ . Suppose, e.g., that  $x_{21} \neq x_{22}$ . Let  $u \in M_1$ ,  $u \notin L_1$ ,  $u \notin L_2$ ,  $N = \{x, u\}^\perp - \{x\}$ . If  $N$  does not contain a point of  $\{y, x_{11}, x_{22}\}^{\perp\perp}$  then by 2.3, and since the point  $x \in \{y, x_{11}, x_{22}\}^\perp$  is collinear with  $u$ , the point  $u$  is collinear with a second point  $x'$  of  $\{y, x_{11}, x_{22}\}^\perp$ . So  $x'$  is collinear with  $y, x_{11}, x_{22}, u$ . Hence  $x' \in \{y, x_{11}, u\}^\perp$  and so  $x' \sim x_{21}$ , giving a triangle  $\{x', x_{21}, x_{22}\}$ , a contradiction. Consequently  $N$  contains a point of

$\{y, x_{11}, x_{22}\}^{\perp\perp}$ . If  $x_{11} \neq x_{12}$ , then an analogous argument shows that  $N$  contains a point of  $M_2$ ; if  $x_{11} = x_{12}$ , then, as  $\{y, x_{11}, x_{22}\}^{\perp\perp} = M_2 \cup \{y\}$ , it is clear that  $N$  contains a point of  $M_2$ . ■

Now we introduce the set  $H_{xy}$ . The elements of  $H_{xy}$  are of two types:

- (a) The sets  $\{x, z\}^\perp - \{y\}$ , with  $x \not\sim z$  and  $y \in \{x, z\}^\perp$ , and
- (b) each set which is the union of all elements of type (b) of  $B_{xy}$  containing a point of some line  $M$  of type (a) of  $B_{xy}$ .

The set  $H_{xy}$  contains  $q^3$  elements of type (a), and by Lemma 3.1.1 it contains  $(q^4 + q^3) : q^2(q^2 - q)/q(q - 1) = q^2 + q$  elements of type (b).

LEMMA 3.1.2. *The elements of  $P_{xy}$  and  $B_{xy}$  in an element of  $H_{xy}$  are the points and lines of a  $2 - (q^2, q, 1)$  design, that is, an affine plane of order  $q$ .*

*Proof.* Easy, using property (G) and Lemma 3.1.1. ■

THEOREM 3.1.3. *The elements of  $P_{xy}$ ,  $B_{xy}$ , and  $H_{xy}$ , respectively, are the points, lines, and planes of affine space  $AG(3, q)$ .*

*Proof.* We have  $|P_{xy}| = q^3$ , each element of  $B_{xy}$  contains  $q$  elements of  $P_{xy}$ , and any two distinct elements of  $P_{xy}$  are contained in a unique element of  $B_{xy}$ . Hence  $\mathcal{S}_{xy}$  is a  $2 - (q^3, q, 1)$  design.

If  $q = 3$ , then by [11, 5.3.2] we have  $\mathcal{S} \cong Q(5, 3)$ . By [11, 3.2.4]  $Q(5, 3) \cong T_3(O)$ , with  $O$  an elliptic quadric of  $PG(3, 3)$ . Now the statement of the theorem is easily checked by taking for  $y$  the point  $(\infty)$  and for  $x$  any point distinct from  $(\infty)$  but collinear with  $(\infty)$ .

Now assume  $q > 3$ . By a theorem of Buekenhout [1] it is sufficient to show that any three points of  $P_{xy}$ , not contained in a common element of  $B_{xy}$ , are contained in a common element of  $H_{xy}$ . Let  $z, u, w$  be three distinct points of  $P_{xy}$ , not in a common element of  $B_{xy}$ , and not in a common element of type (b) of  $H_{xy}$ . Then no two points of  $\{z, u, w\}$  are collinear in  $\mathcal{S}$  (that is, no two points of  $\{z, u, w\}$  are incident with a common line of  $\mathcal{S}$  through  $x$ ), and  $\{y, z, u\}^{\perp\perp}$  has no point in common with  $\{x, w\}^\perp$ . Since  $w \sim x$  and  $x \in \{y, z, u\}^\perp$ , so by 2.3  $w$  is collinear with a second point  $x'$  of  $\{y, z, u\}^\perp$ . Clearly  $\{x, x'\}^\perp$  contains  $y, z, u, w$ , and hence  $z, u, w$  are contained in an element of type (a) of  $H_{xy}$ . ■

COROLLARY 3.1.4. *If  $\mathcal{S} = (P, B, I)$  is a GQ of order  $(q, q^2)$ ,  $q \neq 1$ , for which at least one pair of points  $\{x, y\}$ ,  $x \sim y$ , has property (G), then  $q$  is a prime power.*

*Proof.* Immediate from Theorem 3.1.3. ■

**THEOREM 3.1.5.** *Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(q, q^2)$ ,  $q$  even, for which at least one pair of points  $\{x, y\}$ ,  $x \sim y$ , has property (G). By 2.3 each element of type (a) of  $B_{xy}$  defines a subquadrangle  $\mathcal{S}'$  of order  $q$  of  $\mathcal{S}$ . In this way there arise  $q^3 + q^2$  subquadrangles of order  $q$  of  $\mathcal{S}$ . Also, in any such subquadrangle each pair  $\{y, u\}$  with  $x \in \{y, u\}^\perp$  (and so each pair  $\{x, z\}$  with  $y \in \{x, z\}^\perp$ ) is regular.*

*Proof.* Assume that the elements  $M_1, M_2$  of type (a) of  $B_{xy}$  define the same subquadrangle  $\mathcal{S}'$  of order  $q$ . Let  $M_i \cup \{y\} = \{y, u_i, z_i\}^{\perp\perp}$ ,  $i = 1, 2$ . Then by 2.3 the pair  $\{y, u_i\}$ ,  $i = 1, 2$ , is regular in  $\mathcal{S}'$  and  $\{y, u_i\}^{\perp\perp} = M_i \cup \{y\}$ ,  $i = 1, 2$ . Since  $\{y, u_i\}^{\perp\perp} = \{v_i, w_i\}^{\perp\perp}$  for any two points  $v_i, w_i \in \{y, u_i\}^{\perp\perp}$ ,  $i = 1, 2$ , we have  $\{y, u_1\}^{\perp\perp} \cap \{y, u_2\}^{\perp\perp} = \{y\}$ . Also, as  $x \in \{y, u_i\}^{\perp\perp}$  with  $i = 1, 2$ , any element of type (b) of  $B_{xy}$  containing a point of  $M_1$  also contains a point of  $M_2$ . Hence  $M_1$  and  $M_2$  are parallel lines in the affine plane of type (b) of  $AG(3, q)$  defined by  $M_1$ .

Conversely, let  $M_1, M_2$  be distinct parallel lines of type (a) of an affine plane of type (b) of  $AG(3, q)$ . Suppose that  $M_i \cup \{y\} = \{y, u_i, z_i\}^{\perp\perp}$ ,  $i = 1, 2$ , where  $u_1 \sim u_2$  and  $z_1 \sim z_2$ . If  $w \in \{y, u_1, z_1\}^{\perp\perp} \cap \{y, u_2, z_2\}^{\perp\perp}$ ,  $w \neq x$ , then there arises a triangle  $\{w, u_1, u_2\}$ , a contradiction. Hence  $\{y, u_1, z_1\}^{\perp\perp} \cap \{y, u_2, z_2\}^{\perp\perp} = \{x\}$ . Now we show that each point  $r$  of  $\{y, u_2, z_2\}^{\perp\perp} - \{x\}$  is incident with a line of  $\mathcal{S}$  joining a point of  $\{y, u_1, z_1\}^{\perp\perp}$  to a point of  $\{y, u_1, z_1\}^\perp$ . Suppose the contrary. By 2.3,  $r$  is collinear with two points  $y, y'$  of  $\{y, u_1, z_1\}^{\perp\perp}$ . Now, let  $\{y''\} = \{y, u_2, z_2\}^{\perp\perp} \cap \{x, y'\}^\perp$ . Then there arises the triangle  $\{y', y'', r\}$ , a contradiction. Consequently  $r$  is incident with a line of  $\mathcal{S}$  joining a point  $w_1$  of  $\{y, u_1, z_1\}^{\perp\perp}$  to a point  $v_1$  of  $\{y, u_1, z_1\}^\perp$ . (If  $w_1 \neq y$ , then there arises a triangle  $\{y, r, v_1\}$ , a contradiction. So  $w_1 = y$  and the line  $ry$  of  $\mathcal{S}$  contains a point  $v_1$  of  $\{y, u_1, z_1\}^\perp$ .) Hence  $r$  is a point of the subquadrangle  $\mathcal{S}'$  of order  $q$  defined by  $M_1$ , and so  $\{y, u_2, z_2\}^\perp$  is contained in  $\mathcal{S}'$ . Since  $M_2$  is a line of the affine plane of type (b) defined by  $M_1$ , also  $M_2$  is contained in  $\mathcal{S}'$ . We conclude that  $\mathcal{S}'$  is also the subquadrangle of order  $q$  defined by  $M_2$ .

From the preceding sections it follows that the elements of type (a) of  $B_{xy}$  define  $(q^4 + q^3)/q = q^3 + q^2$  subquadrangles of order  $q$  of  $\mathcal{S}$ .

Now consider a point  $u$  of the subquadrangle  $\mathcal{S}'$  defined by  $M_1$ , with  $y \not\sim u$  and  $x \in \{y, u\}^\perp$ . As  $\{x, u\}^\perp$  has a point in common with  $M_1$ , the point  $u$  belongs to the affine plane  $\pi$  defined by  $M_1$ . Let  $M_2$  be the line of  $\pi$  parallel to  $M_1$ . Then  $M_2$  also defines the subquadrangle  $\mathcal{S}'$ . It follows that  $\{y, u\}$  is regular in  $\mathcal{S}'$ . ■

**Remark 3.1.6.** Assume  $q$  even or odd. From the second section of the proof of the preceding theorem and by interchanging roles of  $x$  and  $y$  in defining affine planes of type (b), we see that  $M_1$  and  $M_2$  are parallel lines



of type (a) in an affine plane of type (b) of  $AG(3, q)$  iff  $(M_1 \cup \{y\})^\perp - \{x\} = N_1$  and  $(M_2 \cup \{y\})^\perp - \{x\} = N_2$  are parallel lines in the affine plane defined by  $y$  and  $N_1$ . ■

### 3.2. $\mathcal{S}$ Has Property (G) either at the Line $L$ or at the Flag $(x, L)$

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(q, q^2)$ ,  $q$  even, which satisfies property (G) at the flag  $(x, L)$ , with  $L = xy$ . Further, let  $x$  be a center of the triad  $\{y, z, u\}$  and let  $\mathcal{S}'$  be the subquadrangle of order  $q$  containing  $\{y, z, u\}^\perp$  and  $\{y, z, u\}^{\perp\perp}$ .

**THEOREM 3.2.1.** *The point  $x$  is regular in  $\mathcal{S}'$ . If  $\mathcal{S}$  satisfies property (G) at the line  $L$ , then all points of  $L$  are regular in  $\mathcal{S}'$ ,  $L$  is regular in  $\mathcal{S}'$ , and  $L$  is regular in  $\mathcal{S}$ .*

*Proof.* Let  $w$  be a point of  $\mathcal{S}'$ , with  $w \not\sim x$  and  $w \not\sim y$ . Further, let  $r, r', r''$  be distinct points of  $\{x, w\}^{\perp'}$ , with  $r \perp L$ . Then  $\{r, r', r''\}^\perp$  and  $\{r, r', r''\}^{\perp\perp}$  are contained in a subquadrangle  $\mathcal{S}''$  of order  $q$  of  $\mathcal{S}$ . Clearly the lines  $xr, xr', xr'', wr, wr', wr''$  are common lines of  $\mathcal{S}'$  and  $\mathcal{S}''$ . Then by [11, 2.3.1] the common points and lines of  $\mathcal{S}'$  and  $\mathcal{S}''$  form a subquadrangle  $\mathcal{S}' \cap \mathcal{S}''$  of order  $(q, t')$ ,  $t' > 1$ , of  $\mathcal{S}''$ . Now by [11, 2.2.2] we have  $\mathcal{S}' \cap \mathcal{S}'' = \mathcal{S}'$ , and so  $\mathcal{S}' = \mathcal{S}''$ . It follows that  $\{r, r'\}^{\perp'\perp} = \{r, r', r''\}^{\perp\perp}$ ,  $\{r, r'\}^{\perp'} = \{r, r', r''\}^\perp$ , and so  $\{r, r'\}$  and  $\{x, w\}$  are regular in  $\mathcal{S}'$ . Hence  $x$  is regular in  $\mathcal{S}'$ .

Assume now that  $\mathcal{S}$  satisfies property (G) at the line  $L$ . With the notations of the previous section and interchanging the roles of  $x$  and  $y$ , we see that also  $y$  is regular in  $\mathcal{S}'$ . Since  $y$  and  $r$  play the same role in  $\mathcal{S}'$ , also  $r$  is regular in  $\mathcal{S}'$ . It follows immediately that all points of  $L$  are regular in  $\mathcal{S}'$ . Hence also the line  $L$  is regular in  $\mathcal{S}'$  (cf. 2.1). Consider now a line  $M$  of  $\mathcal{S}$ ,  $M \not\sim L$ . Let  $y \sim x' \perp M$ ,  $x \sim y' \perp M$ , and  $x'' \in \{y, y'\}^\perp - \{x, x'\}$ . The subquadrangle  $\mathcal{S}'$  of order  $q$  defined by  $x, x', x''$  contains  $L$  and  $M$ . Since  $L$  is regular in  $\mathcal{S}$ , the span  $\{L, M\}^{\perp'\perp} = \{L, M\}^{\perp\perp}$  has size  $q + 1$ . We conclude that  $L$  is regular in  $\mathcal{S}$ . ■

From now on let  $q$  be odd or even and let  $\mathcal{S}$  satisfy property (G) at the flag  $(x, L)$ . Let  $R$  be  $\{u, u_1, u_2\}^{\perp\perp} = \{u, u_1, u_2, \dots, u_q\}$ , with  $\{u, u_1, u_2\}$  a triad having  $x$  as center and with  $u \perp L$ , and let  $L_i$  be the line incident with  $x$  and the point  $u_i$ ,  $i = 1, 2, \dots, q$ . Further, let  $P_{xR}$  be the set of all points, different from  $x$ , collinear with  $x$  and a point of  $R$ , let  $B_{xR} = \{L, L_1, L_2, \dots, L_q\}$ , and let  $C_{xR}$  be the set having as elements the sets  $\{w, w_1, w_2\}^{\perp\perp}$  with  $w \perp L$ ,  $w_1 \perp L_1$ ,  $w_2 \perp L_2$ , and  $x \notin \{w, w_1, w_2\}$ . Also, let incidence  $I_{xR}$  between elements of  $P_{xR}$  and elements of  $B_{xR}$  be induced by the incidence in  $\mathcal{S}$ , and let incidence  $I_{xR}$  between elements of  $P_{xR}$  and elements of  $C_{xR}$  be containment.

**THEOREM 3.2.2.** *The incidence structure  $\mathcal{L} = (P_{xR}, B_{xR}, C_{xR}, I_{xR})$ , with pointset  $P_{xR}$ , lineset  $B_{xR}$ , circleset  $C_{xR}$ , and incidence  $I_{xR}$ , is a Laguerre plane of order  $q$ . Also, for each point  $y \in L$ ,  $y \neq x$ , the derived or internal affine plane  $\mathcal{L}_y$  of  $\mathcal{L}$  at  $y$  is the affine plane  $AG(2, q)$ ; hence, for  $q$  odd  $\mathcal{L}$  is the classical Laguerre plane, that is, arises from the quadratic cone in  $PG(3, q)$ .*

*Proof.* Clearly  $|P_{xR}| = q^2 + q$ ,  $|B_{xR}| = q + 1$ ,  $|C_{xR}| = q^3$ , each line of  $\mathcal{L}$  is incident with  $q$  points of  $\mathcal{L}$ , and each circle of  $\mathcal{L}$  is incident with  $q + 1$  points of  $\mathcal{L}$ .

Let  $\{w, w_1, w_2\}^{\perp\perp}$  and  $\{w', w_1, w_2\}^{\perp\perp}$  be elements of  $C_{xR}$ ,  $w \in L$ ,  $w' \in L$ ,  $w \neq w'$ ,  $w_1 \in L_1$ ,  $w_2 \in L_2$ . Consider  $w'' \in \{w, w_1, w_2\}^{\perp\perp} - \{w, w_1, w_2\}$  and assume  $\{x, w''\}^{\perp}$  has no point in common with  $\{w', w_1, w_2\}^{\perp\perp}$ . As there are no triangles in  $\mathcal{L}$ ,  $w''$  is collinear with no point of  $\{w', w_1, w_2\}^{\perp\perp}$ . Hence  $w''$  is collinear with two distinct points  $x, r$  of  $\{w', w_1, w_2\}^{\perp}$ . Since  $r \sim w''$ ,  $r \sim w_1$ , and  $r \sim w_2$ , we have  $r \sim w$ . So there arises a triangle with vertices  $w, w', r$ , a contradiction. Consequently  $\{x, w''\}^{\perp}$  has a point in common with  $\{w', w_1, w_2\}^{\perp\perp}$ .

Consider again  $\{w, w_1, w_2\}^{\perp\perp}$ . We show that  $\{w, w_1, w_2\}^{\perp\perp}$  has a point in common with each of  $L, L_1, L_2, \dots, L_q$ . If  $w = u$  we are done by Lemma 3.1.1. If  $w \neq u$ , then consider  $\{u, w_1, w_2\}^{\perp\perp}$ , and apply the previous case to  $\{u, u_1, u_2\}^{\perp\perp}$  and  $\{u, w_1, w_2\}^{\perp\perp}$  and the first section of this proof to  $\{u, w_1, w_2\}^{\perp\perp}$  and  $\{w, w_1, w_2\}^{\perp\perp}$ .

It follows that each line of  $\mathcal{L}$  and each circle of  $\mathcal{L}$  have exactly one point in common.

Since  $\mathcal{L}$  satisfies property (G) at  $(x, L)$ , each circle of  $\mathcal{L}$  is uniquely defined by any three of its points; hence two distinct circles of  $\mathcal{L}$  have at most two points in common. Since each triad of points of  $\mathcal{L}$  is on at most one circle of  $\mathcal{L}$ , there are at most  $[(q^2 + q)q^2(q^2 - q)] / [(q + 1)q(q - 1)] = q^3$  circles. As there are exactly  $q^3$  circles, we conclude that each triad of points of  $\mathcal{L}$  is on exactly one circle of  $\mathcal{L}$ .

Consequently  $\mathcal{L}$  is a Laguerre plane of order  $q$ .

Let  $y \in L$ ,  $y \neq x$ . Then the derived or internal affine plane  $\mathcal{L}_y$  of  $\mathcal{L}$  at  $y$  is the affine plane in the statement of Lemma 3.1.2. Now by Theorem 3.1.3 this affine plane  $\mathcal{L}_y$  is  $AG(2, q)$ . Finally, if  $q$  is odd, then by [10]  $\mathcal{L}$  is the classical Laguerre plane, that is, the Laguerre plane arising from the quadratic cone of  $PG(3, q)$ . ■

Let  $O$  be an oval (that is, a set of  $q + 1$  points no three of which are collinear) in a plane  $PG(2, q)$  of some  $PG(3, q)$ , and let  $x$  be a point of  $PG(3, q)$  not contained in  $PG(2, q)$ . Further, let  $K$  be the cone projecting  $O$  from  $x$ . Then the points of  $K - \{x\}$ , the lines on  $K$ , and the intersections of  $K$  by planes not containing  $x$ , are the points, lines, and circles of a Laguerre plane denoted  $\mathcal{L}(O)$  (the incidence is the natural one). If  $q$  is

odd, then  $O$  is necessarily an irreducible conic (see 8.2.4 of [4]) and so  $\mathcal{L}(O)$  is the Laguerre plane arising from the quadratic cone; that is,  $\mathcal{L}(O)$  is the classical Laguerre plane. Further, each known Laguerre plane is isomorphic to a  $\mathcal{L}(O)$ , for some oval  $O$  of some  $PG(2, q)$ .

**THEOREM 3.2.3.** *Suppose that  $q$  is even and that the Laguerre plane  $\mathcal{L}$  of Theorem 3.2.2 is isomorphic to some  $\mathcal{L}(O)$ . Let  $C$  be a circle of  $\mathcal{L}$  and let  $\mathcal{S}'$  be the subquadrangle of order  $q$  containing  $C$  and  $C^\perp$ . Then the projective plane  $\pi_x$  defined by the regular point  $x$  of  $\mathcal{S}'$  is Desarguesian.*

*Proof.* So we assume that  $q$  is even and that  $\mathcal{L} \cong \mathcal{L}(O)$  for some oval  $O$  of some plane  $PG(2, q)$  of  $PG(3, q)$ . The lines of  $\pi_x$  not containing  $x$  are circles of  $\mathcal{L}$  (cf. 2.1 and 3.2). Any two of these lines have exactly one point in common, so we have  $q^2$  circles of  $\mathcal{L}$  which are mutually tangent. Let  $C_1, C_2, \dots, C_{q^2}$  be the corresponding circles of  $\mathcal{L}(O)$ . Since  $q$  is even, the  $q+1$  tangent planes of the cone  $K$  defining  $\mathcal{L}(O)$  contain a common line, the nucleus line  $N$  of  $K$ . The nucleus of the oval  $C_i$  is the intersection of  $N$  with any tangent line of  $C_i$ . Since any two ovals of  $\{C_1, C_2, \dots, C_{q^2}\}$  are mutually tangent, they have a common tangent line, hence have the same nucleus. Consequently  $C_1, C_2, \dots, C_{q^2}$  have a common nucleus  $n \in N$ . Projecting  $K$  minus its vertex  $\bar{x}$  from  $n$ , we see that the incidence structure formed by the points of  $K - \{\bar{x}\}$  and the circles  $C_1, C_2, \dots, C_{q^2}$  is isomorphic to the dual affine plane with as points the lines of  $PG(3, q)$  containing  $n$  and a point of  $K - \{\bar{x}\}$ , and with as lines the planes of  $PG(3, q)$  containing  $n$  but not  $\bar{x}$ . Clearly this dual affine plane is Desarguesian. Hence the dual affine plane obtained from the projective plane  $\pi_x$  by deleting  $x$  and the lines through  $x$  is Desarguesian. We conclude that the projective plane  $\pi_x$  is Desarguesian. ■

#### 4. TRANSLATION GENERALIZED QUADRANGLES OF ORDER ( $s, s^2$ ), $s \neq 1$ , SATISFYING PROPERTY (G)

##### 4.1. GQ Arising from Flocks

Let  $F$  be a flock of the quadratic cone  $K$  with vertex  $x$  of  $PG(3, q)$ , that is, a partition of  $K - \{x\}$  into  $q$  disjoint irreducible conics. Then, by [16], with  $F$  there corresponds a GQ  $\mathcal{S}(F)$  of order  $(q^2, q)$ . In [9] it was shown that  $\mathcal{S}(F)$  satisfies property (G) at its point  $(\infty)$ . If  $\mathcal{S}(F)$  satisfies property (G) at some point  $z \in (\infty)^\perp - \{(\infty)\}$ , then by [12],  $\mathcal{S}(F)$  is isomorphic to the classical GQ  $H(3, q^2)$  arising from the nonsingular Hermitian variety  $H$  of  $PG(3, q^2)$ . Also,  $\mathcal{S}(F) \cong H(3, q^2)$  iff all the planes of the conics of  $F$  contain a common line of  $PG(3, q)$ ; see [16]. Using the fact that any  $\mathcal{S}(F)$  is an elation generalized quadrangle (EGQ) (that is, admits a group  $G$

(elation group) of collineations, each fixing the point  $(\infty)$  linewise, which acts regularly on the points not collinear with  $(\infty)$  it is then clear that for any nonclassical  $\mathcal{S}(F)$  the point  $(\infty)$  is fixed by any collineation and that a nonclassical GQ of order  $(q^2, q)$  is a  $\mathcal{S}(F)$  for at most one point  $(\infty)$ .

Now suppose that the nonclassical GQ  $\mathcal{S}(F)$  of order  $(s, t)$ ,  $s = t^2$ , is a translation generalized quadrangle (TGQ) with base line  $L$ ; that is, assume that:

(i) The GQ  $\mathcal{S}(F)$  admits a group  $G$  of collineations, each fixing  $L$  pointwise, which acts regularly on the set of lines not concurrent with  $L$  (that is,  $\mathcal{S}(F)$  is an EGQ with base line  $L$  and elation group  $G$ ).

(ii) The group  $G$  is abelian.

In such a case,  $G - \{1\}$  is the set of all collineations of  $\mathcal{S}(F)$  fixing  $L$  pointwise and having no fixed line  $M$  with  $M \not\sim L$  (see [11, 8.6.4]). The group  $G$  is called the translation group of the TGQ with base line  $L$ . As  $\mathcal{S}(F)$  is not classical, the point  $(\infty)$  is fixed by all elements of  $G$ . Consequently  $(\infty)$  is incident with  $L$ . Also, N. L. Johnson [6] proved that for such an  $\mathcal{S}(F)$ ,  $s$  (or  $t$ ) is necessarily odd.

By [11, Section 8.7], the dual  $\widehat{\mathcal{S}(F)}$  of the GQ  $\mathcal{S}(F)$  is then isomorphic to a GQ of type  $T(n, 2n, q) = T(O)$ . A GQ  $T(n, 2n, q)$  is described as follows.

In  $PG(4n-1, q)$  consider a set  $O(n, 2n, q) = O$  of  $q^{2n} + 1$   $(n-1)$ -dimensional subspaces  $(\infty) = PG^{(0)}(n-1, q), PG^{(1)}(n-1, q), \dots, PG^{(q^{2n})}(n-1, q)$ , every three of which generate a  $PG(3n-1, q)$ , and such that each element  $PG^{(i)}(n-1, q)$  of  $O$  is contained in a  $PG^{(i)}(3n-1, q)$  having no point in common with any  $PG^{(j)}(n-1, q)$ ,  $j \neq i$  and  $i = 0, 1, \dots, q^{2n}$ . It is easy to check that  $PG^{(i)}(3n-1, q)$  is uniquely determined,  $i = 0, 1, \dots, q^{2n}$ . The space  $PG^{(i)}(3n-1, q)$  is called the tangent space of  $O$  at  $PG^{(i)}(n-1, q)$ . Embed  $PG(4n-1, q)$  in a  $PG(4n, q)$ , and define  $T(n, 2n, q) = T(O)$  as follows. Points are of three types:

(i) The points of  $PG(4n, q) - PG(4n-1, q)$ .

(ii) The  $3n$ -dimensional subspaces of  $PG(4n, q)$  which intersect  $PG(4n-1, q)$  in one of the  $PG^{(i)}(3n-1, q)$ .

(iii) The symbol  $(L)$ .

Lines are of two types:

(a) The  $n$ -dimensional subspaces of  $PG(4n, q)$  which intersect  $PG(4n-1, q)$  in a  $PG^{(i)}(n-1, q)$ .

(b) The elements of  $O$ .

Incidence in  $T(O)$  is defined as follows: A point of type (i) is incident only with lines of type (a); here the incidence is that of  $PG(4n, q)$ . A point of

type (ii) is incident with all lines of type (a) contained in it and with the unique element of  $O$  contained in it. The point  $(L)$  is incident with no line of type (a) and with all lines of type (b). This GQ  $T(O)$  has order  $(q^n, q^{2n})$ .

Here the base point  $(\infty)$  of the EGQ  $\mathcal{S}(F)$  corresponds to the line  $PG^{(o)}(n-1, q) = (\infty)$  of type (b) of  $T(O)$ , and the base line  $L$  of the TGQ  $\mathcal{S}(F)$  corresponds to the point  $(L)$  of type (iii).

Consider any GQ  $T(n, 2n, q) = T(O)$ . By [11, 8.7.2] the  $q^{2n} + 1$  tangent spaces of  $O = O(n, 2n, q)$  form an  $O^* = O^*(n, 2n, q)$  in the dual space of  $PG(4n-1, q)$ . So in addition to  $T(O)$  there arises a TGQ  $T(O^*)$ , also denoted  $T^*(O)$ , with the same parameters. The TGQ  $T^*(O)$  is called the translation dual of the TGQ  $T(O)$ .

Each TGQ  $\mathcal{S}$  of order  $(s, s^2)$ ,  $s \neq 1$ , with base point  $x$  has a kernel  $\mathcal{K}$ , which is a field with a multiplicative group isomorphic to the group of all collineations of  $\mathcal{S}$  fixing the point  $x$ , and any given point not collinear with  $x$ , linewise; we have  $|\mathcal{K}| \leq s$  (see [11]). The field  $GF(q)$  is a subfield of  $\mathcal{K}$  iff  $\mathcal{S}$  is of type  $T(n, 2n, q)$  (here  $s = q^n$ ); see [11]. The TGQ  $\mathcal{S}$  is isomorphic to a  $T_3(O)$  of Tits,  $O$  an ovoid of  $PG(3, s)$ , iff  $|\mathcal{K}| = s$ .

Finally, the TGQ  $T(O)$  of order  $(q^n, q^{2n})$  and its translation dual  $T(O^*)$  have isomorphic kernels.

#### 4.2. The Known GQ $\mathcal{S}(F)$ which Are also TGQ

If  $\mathcal{S}(F)$  is the classical GQ  $H(3, q^2)$ , then it is a TGQ with base line  $L$ ,  $L$  any line of  $\mathcal{S}(F)$ . The dual  $\widehat{\mathcal{S}(F)}$  of  $\mathcal{S}(F)$  is isomorphic to  $T_3(O)$ ,  $O$  an elliptic quadric of  $PG(3, q)$ . Hence the kernel  $\mathcal{K}$  is the field  $GF(q)$ . Also,  $\widehat{\mathcal{S}(F)}$  is isomorphic to its translation dual  $\widehat{\mathcal{S}(F)}^*$ .

Let  $K$  be the quadratic cone with equation  $X_0X_1 = X_2^2$  of  $PG(3, q)$ ,  $q$  odd. Then the  $q$  planes  $\pi_t$  with equation  $tX_0 - mt^\sigma X_1 + X_3 = 0$ ,  $t \in GF(q)$ ,  $m$  a given nonsquare of  $GF(q)$ , and  $\sigma$  a given automorphism of  $GF(q)$ , define a flock  $F$  of  $K$ ; see [16]. All the planes  $\pi_t$  contain the exterior point  $(0, 0, 1, 0)$  of  $K$ . This flock  $F$  is linear; that is, all the planes  $\pi_t$  contain a common line, iff  $\sigma = 1$ . Conversely, every nonlinear flock  $F$  of  $K$  for which the planes of the  $q$  conics all contain a common point, is of the type just described; see [16]. The corresponding GQ  $\mathcal{S}(F)$  were first discovered by W. M. Kantor [8]. Any such GQ  $\mathcal{S}(F)$  is a TGQ for some base line, and so the dual  $\widehat{\mathcal{S}(F)}$  of  $\mathcal{S}(F)$  is isomorphic to some  $T(O)$ . The kernel  $\mathcal{K}$  is the fixed field of  $\sigma$ ; see [13]. Also, in [9] it is proved that  $T(O)$  is isomorphic to its translation dual  $T(O^*)$ .

Let  $K$  be again the quadratic cone with equation  $X_0X_1 = X_2^2$ ,  $q = 3^r$ , and  $r > 2$ . Then the  $q$  planes  $\pi_t$  with equation  $tX_0 - (mt + m^{-t}t^9)X_1 + t^3X_2 + X_3 = 0$ ,  $t \in GF(q)$ ,  $m$  a given nonsquare of  $GF(q)$ , define a flock  $F$  of  $K$ ; see [9]. The corresponding GQ  $\mathcal{S}(F)$  is a TGQ for some base line, and so the dual  $\widehat{\mathcal{S}(F)}$  of  $\mathcal{S}(F)$  is isomorphic to some  $T(O)$ . By [13], the kernel  $\mathcal{K}$  is

here  $GF(3)$ . Payne [9] proves that  $T(O)$  is not isomorphic to its translation dual  $T(O^*)$ . Also, he shows that  $T(O^*)$  is an EGQ which does not arise from a flock. These GQ were called by Payne the Roman GQ.

#### 4.3. $TGQ$ of Order $(s, s^2)$ , $s \neq 1$ , Satisfying Property (G)

Let  $\mathcal{S} = (P, B, I)$  be a GQ of order  $(s, s^2)$ ,  $s \neq 1$ , having property (G) at a pair of points  $\{x, y\}$ ,  $x I L I y$ . Further assume that  $\mathcal{S}$  is a TGQ with base point  $x$ ,  $x I L$ . As the collineation group of  $\mathcal{S}$  is transitive on the points of  $L$  different from  $x$ , it follows that  $\mathcal{S}$  satisfies property (G) at the flag  $(x, L)$ . The GQ  $\mathcal{S}$  is isomorphic to a GQ  $T(O)$ ,  $O = \{\zeta, \zeta_1, \dots, \zeta_{q^{2n}}\}$  with  $q^n = s$ . The point  $x$  corresponds to the point  $(\infty)$  of type (iii) of  $T(O)$ , and we assume that  $L$  corresponds to the line  $\zeta$  of type (b) of  $T(O)$ . The tangent spaces of  $O$  at the elements  $\zeta, \zeta_1, \dots$ , are respectively denoted by  $\tau, \tau_1, \dots$ .

**THEOREM 4.3.1.** *A GQ  $T(O)$  satisfies property (G) at  $\{(\infty), \bar{\zeta}\}$  [or, equivalently, at the flag  $((\infty), \zeta)$ ], with  $\bar{\zeta}$  a point of type (ii) incident with the line  $\zeta$  of type (b) iff for any two elements  $\zeta_i, \zeta_j$  ( $i \neq j$ ) of  $O - \{\zeta\}$  the  $(n-1)$ -dimensional space  $PG(n-1, q) = \tau \cap \tau_i \cap \tau_j$  of  $PG(4n-1, q)$  is contained in exactly  $q^n + 1$  tangent spaces of  $O$ .*

*Proof.* Suppose that  $T(O)$  satisfies property (G) at  $((\infty), \zeta)$ . Let  $\zeta_i, \zeta_j$  be distinct elements of  $O - \{\zeta\}$  and put  $PG(n-1, q) = \tau \cap \tau_i \cap \tau_j$ . By [11, Section 8.7] each point of  $PG(n-1, q)$  is contained in exactly  $q^n + 1$  tangent spaces of  $O$ . Let  $u \in PG(n-1, q)$ , let  $N$  be a line of  $PG(4n, q)$  through  $u$  not contained in  $PG(4n-1, q)$ , and let  $u_1, u_2$  be distinct points of  $N$  not contained in  $PG(4n-1, q)$ . Then each of the points  $N\tau, N\tau_i, N\tau_j$  of type (ii) of  $T(O)$  is collinear with each of the points  $(\infty), u_1, u_2$  of  $T(O)$ . Then  $\{(\infty), u_1, u_2\}^\perp$  is the set  $\{N\tau, N\tau_i, N\tau_j, \dots\}$  with  $\tau, \tau_i, \tau_j, \dots$  the  $q^n + 1$  tangent spaces of  $O$  through  $u$ . As  $T(O)$  satisfies property (G) at  $\zeta$ , we have  $|\{(\infty), u_1, u_2\}^{\perp\perp}| = q^n + 1$ . Hence  $N\tau \cap N\tau_i \cap N\tau_j \cap \dots = \eta$  contains  $q^n$  points of  $PG(4n, q) - PG(4n-1, q)$ . Consequently  $\eta$  is an  $n$ -dimensional space. That is,  $\tau \cap \tau_i \cap \tau_j \cap \dots$  is  $(n-1)$ -dimensional; that is,  $\tau \cap \tau_i \cap \tau_j \cap \dots$  is  $PG(n-1, q)$ . We conclude that  $PG(n-1, q)$  is contained in exactly  $q^n + 1$  tangent spaces of  $O$ .

Conversely, let  $\zeta_i, \zeta_j$  be any two distinct elements of  $O - \{\zeta\}$ , put  $PG(n-1, q) = \tau \cap \tau_i \cap \tau_j$ , and assume that  $PG(n-1, q)$  is always contained in exactly  $q^n + 1$  tangent spaces of  $O$ . Let  $\bar{\zeta}, \bar{\zeta}_i, \bar{\zeta}_j$  be distinct points of type (ii) collinear with  $(\infty)$ , where  $\zeta \subset \bar{\zeta}$ ,  $\zeta_i \subset \bar{\zeta}_i$ ,  $\zeta_j \subset \bar{\zeta}_j$ . Then  $\bar{\zeta} \cap \bar{\zeta}_i \cap \bar{\zeta}_j$  is an  $n$ -dimensional space  $PG(n, q)$ . The  $(n-1)$ -dimensional space  $PG(n-1, q) = PG(n, q) \cap PG(4n-1, q) = \tau \cap \tau_i \cap \tau_j$  is contained in exactly  $q^n + 1$  tangent spaces  $\tau, \tau_i, \tau_j, \dots$  of  $O$ . The  $q^n + 1$  spaces  $PG(n, q)\tau = \bar{\zeta}$ ,  $PG(n, q)\tau_i = \bar{\zeta}_i$ ,  $PG(n, q)\tau_j = \bar{\zeta}_j, \dots$  are points of type (ii) of  $T(O)$ , each of which is collinear with  $(\infty)$  and with the  $q^n$  points of type (i) in  $PG(n, q)$ .

Hence the triad  $\{\bar{\zeta}, \bar{\zeta}_i, \bar{\zeta}_j\}$  is 3-regular. We conclude that  $\mathcal{S}$  satisfies property (G) at  $((\infty), \zeta)$ . ■

**THEOREM 4.3.2.** *If  $q$  is even and if the TGQ  $T(O)$  satisfies property (G) at the flag  $((\infty), \zeta)$ , then the translation dual  $T(O^*)$  satisfies property (G) at the flag  $((\infty), \tau)$ , with  $\tau$  the tangent space of  $O$  at  $\zeta$ .*

*Proof.* Suppose that  $T(O)$  satisfies property (G) at the flag  $((\infty), \zeta)$ , with  $q$  even. Let  $\bar{\zeta}, \bar{\zeta}_i, \bar{\zeta}_j$  be distinct points of type (ii) collinear with  $(\infty)$ , where  $\zeta \subset \bar{\zeta}, \zeta_i \subset \bar{\zeta}_i, \zeta_j \subset \bar{\zeta}_j$ . Then  $\{\bar{\zeta}, \bar{\zeta}_i, \bar{\zeta}_j\}^\perp$  and  $\{\bar{\zeta}, \bar{\zeta}_i, \bar{\zeta}_j\}^{\perp\perp}$  are contained in a subquadrangle  $\mathcal{S}'$  of order  $q^n$  of  $T(O)$ . The space  $\bar{\zeta} \cap \bar{\zeta}_i \cap \bar{\zeta}_j$  is an  $n$ -dimensional space  $PG(n, q)$ , and the space  $PG(n-1, q) = PG(n, q) \cap PG(4n-1, q) = \tau \cap \tau_i \cap \tau_j$  is contained in exactly  $q^n + 1$  tangent spaces  $\tau, \tau_i, \tau_j, \dots$  of  $O$ . Then the points of  $PG(4n, q) - PG(4n-1, q)$  in  $PG(n, q)\zeta, PG(n, q)\zeta_i, PG(n, q)\zeta_j, \dots$  are the points of type (i) in  $\mathcal{S}'$ . Now we consider a line of type (a) concurrent with  $\zeta$  and incident with a point of type (i) in the space  $PG(n, q)\zeta_i$ . As this line is also a line of  $\mathcal{S}'$ , it is incident with a point of type (i) in each of the spaces  $PG(n, q)\zeta_j, \dots$ . Hence the spaces  $PG(n, q)\zeta_j, \dots$  are contained in the  $3n$ -dimensional space  $PG(n, q)\zeta\zeta_i = PG(3n, q)$ . Consequently the spaces  $PG(n-1, q), \zeta, \zeta_i, \zeta_j, \dots$  are contained in a common  $PG(3n-1, q) = PG(3n, q) \cap PG(4n-1, q)$ . So, by [11, 8.7.2], the  $(3n-1)$ -dimensional space  $\zeta\zeta_i\zeta_j$  contains exactly  $q^n + 1$  elements of  $O$ . Applying Theorem 4.3.1 to  $O^*$  now gives that the translation dual  $T(O^*)$  of  $T(O)$  has property (G) at the flag  $((\infty), \tau)$ . ■

**Remark 4.3.3.** From the proof of Theorem 4.3.2 follows that for  $q$  even the  $(n-1)$ -dimensional space  $\tau \cap \tau_i \cap \tau_j$  is contained in the  $(3n-1)$ -dimensional space  $\zeta\zeta_i\zeta_j$ .

Assume again that the TGQ  $T(O)$  has property (G) at the flag  $((\infty), \zeta)$ . We consider now the translation dual  $T(O^*)$  of  $T(O)$ . We have  $O^* = \{\tau, \tau_1, \dots, \tau_{q^{2n}}\}$ ,  $\zeta$  is the tangent space of  $O^*$  at  $\tau$ , and  $\zeta_i$  is the tangent space of  $O^*$  at  $\tau_i$ ,  $i = 1, 2, \dots, q^{2n}$ . Also, in the space  $PG(4n-1, q)$  of  $O^*$  the  $(3n-1)$ -dimensional space  $PG(3n-1, q) = \tau\tau_i\tau_j$  ( $i \neq j$ ), with  $\tau_i$  and  $\tau_j$  any two distinct elements of  $O^*$ , contains exactly  $q^n + 1$  elements of  $O^*$ . In the  $PG(4n, q)$  containing  $T(O^*)$  we now consider a  $PG(3n, q)$  containing  $PG(3n-1, q)$  but not contained in  $PG(4n-1, q)$ . Then the point  $(\infty)$  together with the points and lines of  $T(O^*)$  in  $PG(3n, q)$  define a TGQ  $T(\tilde{O}^*)$  of order  $s = q^n$ , where  $\tilde{O}^*$  is the set of all elements of  $O^*$  in  $PG(3n-1, q)$  (see [11, Section 8.7]).

**THEOREM 4.3.4.** *The TGQ  $T(O^*)$  contains at least  $s^3 + s^2$  subquadrangles (TGQ)  $T(\tilde{O}^*)$  of order  $s$ . The projective plane  $\pi_\tau$  of order  $s$  defined by the regular line  $\tau$  of  $T(\tilde{O}^*)$  is Desarguesian. It follows that for  $s$  odd, each of the  $s^3 + s^2$  GQ  $T(\tilde{O}^*)$  is isomorphic to the classical GQ  $Q(4, s)$ .*

*Proof.* Project the elements of  $O^* - \{\tau\}$  from  $\tau$  onto a  $(3n-1)$ -dimensional space  $\eta$  skew to  $\tau$ . The projection of  $\tau_i$  from  $\tau$  onto  $\eta$  is denoted by  $\tau'_i$ ,  $i = 1, 2, \dots, q^{2n}$ . Then  $\tau'_1 \cup \tau'_2 \cup \dots \cup \tau'_{q^{2n}}$  is  $\eta - \zeta'$ , where  $\zeta'$  is the  $(2n-1)$ -dimensional space  $\zeta \cap \eta$ . Further, the set of all intersections  $\tau\tau_i\tau_j \cap \eta$ ,  $i \neq j$ , is denoted by  $\mathcal{B}$ . Then with respect to inclusion the elements of  $\mathcal{P} = \{\tau'_1, \dots, \tau'_{q^{2n}}\}$  and  $\mathcal{B}$ , are the points and lines, respectively, of a  $2 - (q^{2n}, q^n, 1)$  design  $\mathcal{A}$ , that is, of an affine plane of order  $q^n = s$ . If  $q'$  and  $q''$  are distinct parallel lines of  $\mathcal{A}$ , then  $q' \cap q''$  is necessarily an  $(n-1)$ -dimensional subspace of  $\zeta'$ . Hence all lines of  $\mathcal{A}$  parallel to  $q'$  contain a common  $(n-1)$ -dimensional subspace of  $\zeta'$ . These  $q^n + 1$  (distinct) subspaces of  $\zeta'$  are denoted by  $v_0, v_1, \dots, v_{q^n}$ . If for some  $i \neq j$   $v_i \cap v_j \neq \emptyset$ , say  $x \in v_i \cap v_j$ , then lines  $q'$  and  $q''$  of  $\mathcal{A}$  containing respectively  $v_i$  and  $v_j$ , do not intersect in a point of  $\mathcal{A}$ , so are parallel, that is  $v_i = v_j$ , a contradiction. Hence  $\{v_0, v_1, \dots, v_{q^n}\}$  is a partition (a spread) of  $\zeta'$ . It is clear that the elements  $v_0, v_1, \dots, v_{q^n}$  and  $\zeta'$  extend  $\mathcal{A}$  to a projective plane  $\Pi$  of order  $q^n$ . Now by a theorem of Segre [14] such a plane  $\Pi$  is always Desarguesian; that is, in  $\eta$  there are  $n$  planes  $\pi_1, \pi_2, \dots, \pi_n$  over  $GF(q^n)$ , which are conjugate with respect to the  $n$ th extension  $GF(q^n)$  of  $GF(q)$  and which generate  $\eta$ , such that each plane  $\pi_i$  has a point in common with each element of the pointset of  $\Pi$  (any point of  $\Pi$  is generated by a point of  $\pi_1$  and its conjugates).

Now we consider the projective plane  $\pi_\tau$  of order  $s = q^n$  defined by the regular line  $\tau$  of  $T(\tilde{O}^*)$ . Let  $\tilde{O}^*$  be defined by  $PG(3n-1, q) = \tau\tau_i\tau_j$  and let  $T(\tilde{O}^*)$  be contained in  $PG(3n, q)$ . Further, let  $\tau\tau_i\tau_j \cap \eta = \bar{\eta}$  and let  $\bar{\eta} \subset PG(2n, q) \subset PG(3n, q)$ , with  $PG(2n, q) \not\subset \tau\tau_i\tau_j$ . The points of  $\Pi$  in  $\bar{\eta}$  form the spread  $S$ . Identifying each line of type (a) of  $T(\tilde{O}^*)$  concurrent with  $\tau$  with its intersection with  $PG(2n, q)$ , each line of type (b) of  $T(\tilde{O}^*)$  distinct from  $\tau$  with its projection from  $\tau$  onto  $\bar{\eta}$ , and  $\tau$  with  $\zeta \cap \bar{\eta}$ , we see that  $\pi_\tau$  is isomorphic to the projective translation plane  $\pi(S)$  defined by the spread  $S$  (cf. [3]). By the end of the previous section,  $S$  is a regular  $(n-1)$ -spread (see [5, Sect. 25.6]) of  $\bar{\eta}$ . Hence  $\pi(S)$ , and so  $\pi_\tau$ , is Desarguesian (see [5, Sect. 25.7]).

Finally, suppose that  $s$  is odd. As the spread  $S$  is regular, it follows from a result by Casse, Thas, and Wild (see [11, 8.7.7]) that  $T(\tilde{O}^*)$  is isomorphic to the classical GQ  $Q(4, s)$  arising from the nonsingular quadric of  $PG(4, s)$ . ■

**LEMMA 4.3.5.** *Suppose that  $T(\tilde{O}^*)$  is isomorphic to the classical GQ  $Q(4, s)$ , with  $\tilde{O}^*$  being contained in the space  $PG(3n-1, q)$ . Then  $\tilde{O}^*$  can be extended to an  $(n-1)$ -spread  $\tilde{S}$  of  $PG(3n-1, q)$  having the property that any  $(2n-1)$ -dimensional space generated by two distinct elements of  $\tilde{S}$  contains exactly  $q^n + 1$  elements of  $\tilde{S}$ . These  $(2n-1)$ -dimensional spaces*



together with the elements of  $\tilde{S}$ , and with containment as incidence, are the lines and points of a Desarguesian plane  $PG(2, q^n)$ .

*Proof.* The last part of the statement follows from a theorem of Segre [14]. The first part easily follows from:

(a) Let  $x \sim (\infty) \sim y$ ,  $x \not\sim y$ . If  $u, v \in \{x, y\}^\perp$  with  $u \neq (\infty) \neq v$ ,  $(\infty) \not\sim L$  with  $x \not\sim L$ ,  $y \not\sim L$ , and  $u \not\sim U \sim L$ ,  $v \not\sim V \sim L$ , then each point of  $\{x, y\}^\perp$  is incident with a line of  $\{U, V\}^\perp$  (or, equivalently  $\{U, V\}^{\perp\perp}$ ).

(b) Consider  $\{L_1, N_1\}^\perp$ ,  $\{L_2, N_2\}^\perp$ , with  $(\infty) \not\sim L_1$ ,  $(\infty) \not\sim L_2$ ,  $(\infty) \not\sim N_1 \not\sim L_1$ ,  $(\infty) \not\sim N_2 \not\sim L_2$ ,  $L_2 \notin \{L_1, N_1\}^{\perp\perp} \cup \{L_1, N_1\}^\perp$ . Then the lines of  $\{L_1, N_1\}^\perp$  and  $\{L_2, N_2\}^\perp$  are incident with  $q^n + 1$  common points. The set of these points is denoted by  $D$ . If  $u, v$  are two distinct points of  $D$ , with  $u \neq (\infty) \neq v$ , if  $(\infty) \not\sim L$  with  $L \notin \{L_i, N_i\}^{\perp\perp} \cup \{L_i, N_i\}^\perp$ ,  $i = 1, 2$ , and if  $u \not\sim U \sim L$ ,  $v \not\sim V \sim L$ , then each point of  $D$  is incident with a line of  $\{U, V\}^\perp$  (or, equivalently  $\{U, V\}^{\perp\perp}$ ).

(c) From (a) and (b) follows that  $\tilde{O}^*$  can be extended to an  $(n-1)$ -spread  $\tilde{S}$  of  $PG(3n-1, q)$ , that each conic of  $Q$  through  $(\infty)$  is in  $T_2(\tilde{O}^*)$  of the form  $\{(\infty)\} \cup (PG(n, q) - PG(3n-1, q))$  with  $PG(n, q)$  an  $n$ -dimensional subspace of  $PG(3n, q)$  intersecting  $PG(3n-1, q)$  in an element of  $\tilde{S} - \tilde{O}^*$ , that each  $(2n-1)$ -dimensional subspace of  $PG(3n-1, q)$  containing at least two elements of  $\tilde{O}^*$  contains exactly  $q^n + 1$  elements of  $\tilde{S}$ , and that each  $(2n-1)$ -dimensional subspace of  $PG(3n-1, q)$  containing exactly one element of  $\tilde{O}^*$  and being disjoint from all other  $q^n$  elements of  $\tilde{O}^*$  contains exactly  $q^n + 1$  elements of  $\tilde{S}$ .

(d) From (c) follows that each elliptic quadric on  $Q$  through  $(\infty)$  is in  $T_2(\tilde{O}^*)$  of the form  $\{(\infty)\} \cup (PG(2n, q) - PG(3n-1, q))$  with  $PG(2n, q)$  a  $2n$ -dimensional subspace of  $PG(3n, q)$  intersecting  $PG(3n-1, q)$  in a  $(2n-1)$ -dimensional space containing exactly  $q^n + 1$  elements of  $\tilde{S} - \tilde{O}^*$ . The number of these  $(2n-1)$ -dimensional spaces is equal to  $(q^n - 1)q^n/2$ . ■

**COROLLARY 4.3.6.** *The dual  $\widehat{\mathcal{S}(F)}$  of a TGQ  $\mathcal{S}(F)$  of Kantor (cf. 4.2) contains at least  $s^3 + s^2$  classical subquadrangles  $Q(4, s)$ . Equivalently,  $\mathcal{S}(F)$  contains at least  $s^3 + s^2$  classical subquadrangles  $W(s)$ . Any Roman TGQ of Payne (cf. 4.2) contains at least  $s^3 + s^2$  classical subquadrangles  $Q(4, s)$ .*

Let  $\mathcal{S}$  be a GQ of order  $(s, s^2)$ ,  $s$  even, having property (G) at the flag  $(x, L)$  and suppose that  $\mathcal{S}$  is a TGQ with base point  $x$ . Further, let  $x$  be a center of the triad  $\{y, z, u\}$ , with  $y \not\sim L$ , and let  $\mathcal{S}'$  be the subquadrangle of order  $s$  containing  $\{y, z, u\}^\perp$  and  $\{y, z, u\}^{\perp\perp}$ . As  $L$  is a regular line of the TGQ  $\mathcal{S}$ , it is also a regular line of  $\mathcal{S}'$ .

**THEOREM 4.3.7.** *For  $s$  even, the projective plane  $\pi_L$  arising from the regular line  $L$  of  $\mathcal{S}'$  is Desarguesian.*

*Proof.* By Theorem 4.3.2 the translation dual  $T(O^*)$  of  $T(O) \cong \mathcal{S}$  has property (G) at the flag  $((\infty), \tau)$  with  $\tau$  the tangent space of  $O$  at  $\zeta$ . So by Theorem 4.3.1 the  $(3n-1)$ -dimensional space  $PG(3n-1, q) = \zeta\zeta_i\zeta_j$ , with  $\zeta_i$  and  $\zeta_j$  any two distinct elements of  $O - \{\zeta\}$ , contains exactly  $q^n + 1$  elements of  $O$ . If  $\tilde{O}$  is the set of these  $q^n + 1$  elements, then  $\mathcal{S}'$  is of type  $T(\tilde{O})$ , where  $T(\tilde{O})$  is defined by  $\tilde{O}$  and a  $3n$ -dimensional subspace  $PG(3n, q)$  containing  $PG(3n-1, q)$  but not contained in  $PG(4n-1, q)$  (cf. proof of 4.3.2). By Theorem 4.3.4 the projective plane  $\pi_\zeta$  of order  $s$  defined by the regular line  $\zeta$  of  $T(\tilde{O})$  is Desarguesian; hence  $\pi_L$  is Desarguesian. ■

## 5. OVOIDS OF THE GENERALIZED QUADRANGLE $Q(4, s)$ , $s$ ODD

### 5.1. Definition and Known Results

An ovoid  $\mathcal{O}$  of the GQ  $Q(4, s)$ ,  $s$  odd, is a set of  $s^2 + 1$  points, no two of which are collinear; equivalently  $\mathcal{O}$  has a point in common with each line of  $Q(4, s)$ .

If  $Q \subset PG(4, s)$  and if the hyperplane  $PG(3, s)$  intersects  $Q$  in an elliptic quadric  $\mathcal{O}$ , then  $\mathcal{O}$  is an ovoid of  $Q(4, s)$ . For  $s$  an even power of 2 no other ovoids of  $Q(4, s)$  are known. For  $s = 2^{2h+1}$ ,  $h \geq 1$ , one other type of ovoid is known; its projection from the nucleus  $n$  of  $Q$  onto a hyperplane not containing  $n$ , is the Tits ovoid; see [15]. For  $s$  odd, Kantor [7] constructed two types of nonclassical ovoids: (i) type  $\mathcal{K}_1$ , here  $s = p^h$ ,  $p$  an odd prime and  $h > 1$ , and these ovoids are characterized by the property that they are in just one way the union of  $s$  conics which are mutually tangent at a common point (see also [16]), and (ii) type  $\mathcal{K}_2$ , here  $s = 3^{2h+1}$ ,  $h \geq 1$ , and each such ovoid arises from a Ree–Tits ovoid on a nonsingular quadric in projective 6-space.

With each ovoid of  $Q(4, s)$  there corresponds a spread, which is a partition by lines, of the dual GQ  $W(s)$ . And with a spread of  $W(s)$  there corresponds, by the standard argument, a translation plane  $\mathcal{P}$  of order  $s^2$ . If  $\mathcal{O}$  is the elliptic quadric, then  $\mathcal{P}$  is Desarguesian. If  $\mathcal{O}$  is of Tits type, then  $\mathcal{P}$  is the Lüneburg plane; see [15]. If  $\mathcal{O}$  is of Kantor type  $\mathcal{K}_1$ , then  $\mathcal{P}$  is a Knuth semifield plane; see [7].

### 5.2. Subquadrangles and Ovoids

Let  $\mathcal{S}'$  be a subquadrangle of order  $(s, t')$  of the GQ  $\mathcal{S}$  of order  $(s, t)$ ,  $t' < t$ . If  $z$  is a point of  $\mathcal{S}$  which is not contained in  $\mathcal{S}'$ , then the  $1 + st'$  points of  $\mathcal{S}'$  collinear with  $z$  form an ovoid  $\mathcal{O}$  of  $\mathcal{S}'$ ; see [11, 2.2.1]. For example, if  $\mathcal{S} = Q(5, s)$  and  $\mathcal{S}' = Q'(4, s)$ , with  $Q' \subset Q$ , then  $\mathcal{O}$  is an elliptic quadric on  $Q'$ .

### 5.3. Ovoids of $T(\tilde{O}^*) \cong Q(4, s)$ Arising from $T(O^*)$

Assume that the TGQ  $T(O)$ , with  $O = \{\zeta, \zeta_1, \dots, \zeta_{q^{2n}}\}$  consisting of  $q^{2n} + 1$   $(n-1)$ -dimensional subspaces in  $PG(4n-1, q)$ , has property (G) at the flag  $((\infty), \zeta)$ . Let  $T(O^*)$  be the translation dual of  $T(O)$ . Then  $O^* = \{\tau, \tau_1, \dots, \tau_{q^{2n}}\}$ , with  $\tau$  the tangent space of  $O$  at  $\zeta$  and  $\tau_i$  the tangent space of  $O$  at  $\zeta_i$ ,  $i = 1, 2, \dots, q^{2n}$ . Further, let  $q$  be odd. Then by Theorem 4.3.4 the GQ  $T(O^*)$  has  $s^3 + s^2$ ,  $s = q^n$ , subquadrangles  $T(\tilde{O}^*)$  isomorphic to  $Q(4, s)$ .

If  $z$  is a point of  $T(O^*)$  which is not contained in the subquadrangle  $T(\tilde{O}^*)$ , then the corresponding ovoid of  $T(\tilde{O}^*)$  will be denoted by  $\mathcal{O}(z)$ .

**THEOREM 5.3.1.** *If  $T(O)$  is a TGQ of Kantor type (including the classical GQ  $Q(5, s)$ ), then  $\mathcal{O}(z)$  is always an elliptic quadric or of Kantor type  $\mathcal{K}_1$ .*

*Proof.* Let  $T(O)$  be a TGQ of Kantor type (cf. 4.2). Then  $T(O) \cong T(O^*)$ . Let  $\tilde{O}^*$  be contained in  $PG(3n-1, q)$  and assume that  $\tilde{O}^* = \{\tau, \tau_1, \dots, \tau_{q^n}\}$ . Then  $\tau_i$  (resp.  $\tau$ ) is contained in exactly one  $(2n-1)$ -dimensional space  $\tilde{\zeta}_i$  (resp.  $\tilde{\zeta}$ ) of  $PG(3n-1, q)$ , having an empty intersection with each element of  $\tilde{O}^* - \{\tilde{\zeta}_i\}$  (resp.  $\tilde{O}^* - \{\tilde{\zeta}\}$ ),  $i = 1, 2, \dots, q^n$ . Since  $T(O^*)$  has property (G) at the flag  $((\infty), \tau)$ , the  $(n-1)$ -spaces  $\tilde{\zeta} \cap \tilde{\zeta}_i$  are exactly the  $(n-1)$ -spaces  $\tilde{\zeta} \cap \tilde{\zeta}_j$ ,  $i = 1, 2, \dots, q^n$ , and  $j = 1, 2, \dots, q^{2n}$ .

First, let  $z$  be a point of type (ii) of  $T(O^*)$  which is not contained in  $T(\tilde{O}^*)$ . Then  $\mathcal{O}(z) = \{(\infty)\} \cup (PG(2n, q) - PG(3n-1, q))$ , where  $PG(2n, q)$  is a subspace of the space  $PG(3n, q)$  in which  $T(\tilde{O}^*)$  is defined and with  $PG(2n, q) \cap PG(3n-1, q)$  of the form  $\tilde{\zeta}_j \cap PG(3n-1, q)$ ,  $j \in \{q^{n+1}, \dots, q^{2n}\}$ . Let us now identify  $T(\tilde{O}^*)$  with the GQ  $T_2(C)$ , where  $C$  is a conic of  $PG(2, s)$ . Then in  $T_2(C)$  the ovoid  $\mathcal{O}(z)$  consists of  $(\infty)$  and the points of type (i) on  $s$  lines of  $PG(3, s)$  containing a common point on a tangent line of  $C$ . So in  $Q(4, s) \cong T_2(C)$ ,  $\mathcal{O}(z)$  consists of  $s$  conics which are mutually tangent at a common point  $(\infty)$ .

Next, let  $z$  be a point of type (i) of  $T(O^*)$  which is not contained in  $T(\tilde{O}^*)$ . Then  $\mathcal{O}(z)$  consists of  $s+1$  points of type (ii) and the  $s^2-s$  points  $u_j$ , with  $\{u_j\} = \tau_j z \cap PG(3n, q)$  and  $j = s+1, \dots, s^2$ . The  $s^2-s$  spaces  $\tau_j$ ,  $j = s+1, \dots, s^2$ , are distributed over  $s-1$   $(3n-1)$ -dimensional spaces containing  $\tilde{\zeta}$ . Hence the  $s^2-s$  points  $u_j$  are distributed over  $s-1$   $2n$ -dimensional spaces  $\pi_1, \dots, \pi_{s-1}$  containing  $\tilde{\zeta}$ . If  $u_j$  and  $u_{j'}$ ,  $j \neq j'$ , belong to a common space  $\pi_k$ , then  $u_j u_{j'}$  is skew to  $\tau$  and  $\tau_j \tau_{j'} \cap \tilde{\zeta}$  is one of the spaces  $\tilde{\zeta} \cap \tilde{\zeta}_i$ ,  $i = 1, 2, \dots, q^n$  (cf. Lemma 4.3.5). It follows that for any three distinct points  $u_j$ ,  $u_{j'}$ , and  $u_{j''}$  of a common space  $\pi_k$ , the points  $u_j u_{j'} \cap \tilde{\zeta}$ ,  $u_j u_{j''} \cap \tilde{\zeta}$ ,  $u_{j'} u_{j''} \cap \tilde{\zeta}$  lie in different spaces  $\tilde{\zeta} \cap \tilde{\zeta}_i$ . In  $T_2(C)$  the ovoid  $\mathcal{O}(z)$  consists of  $s+1$  points of type (ii) and  $s^2-s$  points  $u_j$  of type (i) distributed over  $s-1$  planes  $\pi_1, \dots, \pi_{s-1}$  containing a common tangent line  $Z$  of  $C$ . The common point of  $Z$  and  $C$  is denoted by  $t$ . By a preceding argument the point  $t$  together with the  $s$  points  $u_j$  in a common plane  $\pi_k$  constitute a set of  $s+1$

points no three of which are collinear, that is, an oval of  $\pi_k$ . As  $s$  is odd this oval is a conic  $C_k$  (see 8.2.4 of [4]). Clearly  $Z$  is the tangent line of  $C_k$  at  $t$ . In  $T_2(C)$  all points of  $C_k - \{t\}$  are collinear with the point  $\pi_k$  of type (ii). Now it is clear that in  $Q(4, s) \cong T_2(C)$   $\mathcal{O}(z)$  consists of  $s+1$  points collinear with  $(\infty)$  and of the  $s^2-s$  points of  $s-1$  sets  $\bar{C}_k$ , where  $C'_k = \bar{C}_k \cup \{z_k\}$  is a conic and the points  $z_1, z_2, \dots, z_{s-1}$  are on a line  $T$  through  $(\infty)$ . All points of  $C'_k$  are collinear with a point  $w_k$  on  $T$ , and hence also with a second point  $w'_k$ . If  $z_1 \notin \mathcal{O}(z)$ , then, as each line through  $z_1$  contains exactly one point of  $\mathcal{O}(z)$ , the point  $z_1$  is necessarily one of the points  $w_k$ , say  $w_2$ . As  $v \in \bar{C}_2$  is collinear with no point of  $\bar{C}_1$ , the line  $z_1 v$  contains the point  $w'_1$ . Hence  $w'_1$  is on each line  $z_1 v$ , a contradiction. Consequently  $z_1 = z_2 = \dots = z_{s-1} = \bar{z}$  is the point of  $\mathcal{O}(z)$  on the line  $T$ . As  $w'_1$  is collinear with no point of  $\bar{C}_k$ ,  $k \neq 1$ , we necessarily have  $w'_1 \sim w'_k$ . It follows that  $w'_1, w'_2, \dots, w'_{s-1}$  are on a common line  $\bar{W}$  through  $\bar{z}$ . Hence the planes of the conics  $C'_1, C'_2$  are in the polar space  $\eta$  of  $w_1 w'_1 \cap w_2 w'_2$  with respect to  $Q$ . So  $C'_1, C'_2$  are on the elliptic quadric  $\eta \cap Q$ , and since  $C'_1 \cap C'_2 = \{\bar{z}\}$  they are tangent at  $\bar{z}$ . Consequently  $C'_1, \dots, C'_{s-1}$  are mutually tangent at  $\bar{z}$ . Clearly no two of the points  $\bar{z}, w'_1, \dots, w'_{s-1}$  coincide. Let  $w'_s$  be the remaining point of the line  $\bar{W}$ . No point of  $\bar{C}_1 \cup \bar{C}_2 \cup \dots \cup \bar{C}_{s-1}$  is collinear with  $w'_s$ , and so  $w'_s$  is collinear with the  $s+1$  points of  $\mathcal{O}(z)$  in  $(\infty)^\perp$ . It follows that the set  $C'_s$  of all points of  $\mathcal{O}(z)$  collinear with  $(\infty)$  is the conic  $\{(\infty), w'_s\}^\perp$ . Now an argument used before shows that  $C'_s$  and  $C'_k$ ,  $k=1, 2, \dots, s-1$ , are mutually tangent at  $\bar{z}$ . Consequently the conics  $C'_1, C'_2, \dots, C'_s$  are mutually tangent at  $\bar{z}$ ; that is,  $\mathcal{O}(z)$  is an elliptic quadric or of Kantor type  $\mathcal{K}_1$ . ■

*Conjecture 5.3.2.* If  $T(O^*)$  is a Roman GQ and  $q=3^r$  with  $r>2$ , then for each point  $z$  of type (i) not contained in  $T(\bar{O}^*)$ , the ovoid  $\mathcal{O}(z)$  is neither an elliptic quadric nor of Kantor type  $\mathcal{K}_1$ . Then it is clear that for  $r$  even, the ovoid  $\mathcal{O}(z)$  is new.

*Note added in proof.* This conjecture is proved in: J. A. THAS AND S. E. PAYNE, Spreads and ovoids in finite generalized quadrangles, *Geom. Dedicata*, to appear. For any  $r \geq 3$  the avoid, and also the corresponding plane, are new.

*Part II.* The goal of Part II is the classification of all translation generalized quadrangles of order  $(s, s^2)$ ,  $s > 1$ , satisfying property (G) at some flag  $((\infty), L)$ . At this moment many partial results are already obtained.

#### ACKNOWLEDGMENT

I thank my colleague and friend S. E. Payne for many interesting discussions on the subject.

## REFERENCES

1. F. BUEKENHOUT, Une caractérisation des espaces affins basée sur la notion de droite, *Math. Z.* **111** (1969), 367–371.
2. P. J. CAMERON AND J. H. VAN LINT, “Designs, Graphs, Codes and Their Links,” Cambridge Univ. Press, Cambridge, 1991.
3. P. DEMBOWSKI, “Finite Geometries,” Springer, Berlin, 1968.
4. J. W. P. HIRSCHFELD, “Projective Geometries over Finite Fields,” Oxford Univ. Press, Oxford, 1979.
5. J. W. P. HIRSCHFELD AND J. A. THAS, “General Galois Geometries,” Oxford Univ. Press, Oxford, 1991.
6. N. L. JOHNSON, Semifield flocks of quadratic cones, *Simon Stevin* **61** (1987), 313–326.
7. W. M. KANTOR, Ovoids and translation planes, *Canad. J. Math.* **34** (1982), 1195–1203.
8. W. M. KANTOR, Some generalized quadrangles with parameters  $(q^2, q)$ , *Math. Z.* **192** (1986), 45–50.
9. S. E. PAYNE, An essay on skew translation generalized quadrangles, *Geom. Dedicata* **32** (1989), 93–118.
10. S. E. PAYNE AND J. A. THAS, Generalized quadrangles with symmetry, II, *Simon Stevin* **49** (1976), 81–103.
11. S. E. PAYNE AND J. A. THAS, “Finite Generalized Quadrangles,” Pitman, London, 1984.
12. S. E. PAYNE AND J. A. THAS, Conical flocks, partial flocks, derivation, and generalized quadrangles, *Geom. Dedicata* **38** (1991), 229–243.
13. L. A. ROGERS, Characterization of the kernel of certain translation generalized quadrangles, *Simon Stevin* **64** (1990), 319–328.
14. B. SEGRE, Teoria di Galois, fibrazioni proiettive e geometrie non desarguesiane, *Ann. Mat. Pura Appl.* **64** (1964), 1–76.
15. J. A. THAS, Ovoidal translation planes, *Arch. Math.* **23** (1972), 110–112.
16. J. A. THAS, Generalized quadrangles and flocks of cones, *European J. Combin.* **8** (1987), 441–452.